

# Palaeozoic volcanic-hosted massive sulphide deposits

J. Bruce Gemmill, Ross R. Large & Khin Zaw

Centre for Ore Deposit Research, University of Tasmania, GPO Box 252-79, Hobart, TAS 7001, Australia.

## EXPLORATION MODEL

### Examples

Eastern Australia: Hellyer, Rosebery, Que River, Hercules, Mt Lyell, Woodlawn, Thalanga, Wilga, Currawong, Balcooma, Mt Chalmers. Other major examples include deposits from the Japanese Green Tuff belt (Kuroko deposits), Norwegian Caledonides and Canadian Bathurst Group, N.B.

### Target

- Major deposit size: 15–90 Mt.
- Median deposit size: 1–5 Mt.
- Average grade for Cu–Pb–Zn deposits: 1.0% Cu, 12% Zn, 5% Pb.
- High Ag and Au credits: average 120 g/t Ag and 2.0 g/t Au.

### Mining and treatment

- Massive sulphide nature limits dilution effect.
- Stringer zones only mined where high-grade or large tonnage (e.g. Mt Lyell).
- Polymetallic nature may cause recovery problems in fine-grained ores.
- Metamorphosed–recrystallised ores are easier to treat.
- Fe content of sphalerite is moderate to high.
- Pyrite content is high.




Figure 1. Cross-section of typical mound-style deposit.




Figure 2. Regional schematic (based on Mt Windsor Province, Queensland).

### Regional geological criteria

- Back-arc and inter-arc rift volcanic basins.
- Preferred ages: Cambro-Ordovician and Silurian.
- Calc-alkaline submarine volcanics and sediments.
- Compositional variation: rhyolite–andesite–basalt.
- Proximity to syn-volcanic rift faults.
- Located proximal to volcanic centres (Cu-rich ores) or in distal volcanic facies (Pb–Zn-rich ores).
- Syn-volcanic magnetite-series granites may be present.
- Rhyolite is most common footwall composition.
- Sediment and/or mafic volcanics are most common hangingwall rock types.
- Regional sericite ± chlorite alteration in footwall volcanics.

### Local geological criteria

- Ore located in favourable horizon between volcanic units.
- Favourable horizon may be iron-rich exhalite, sulphide-bearing epiclastic, shale or carbonate.
- Ore same age as host volcanic–sedimentary rocks.
- Deposits vary in shape from blankets to lenses, to mounds and pipes.
- Large (1992) recognises ten different styles of VHMS deposits.

### Mineralisation features

- Zn–Pb massive sulphide lens is stratiform.
- Massive sulphide may be banded, brecciated or massive and featureless.
- Cu-rich footwall stringer zone is cross-cutting.
- Chlorite, sericite, quartz, barite, carbonate are major gangue minerals.
- Vertical (up-stratigraphy) zonation of Cu, Au→Pb, Zn, Ag, Au→Ba.
- Pyrite is major sulphide mineral plus sphalerite, galena, chalcocopyrite ± tetrahedrite, arsenopyrite.
- Magnetite and pyrrhotite are rare except for strongly metamorphosed deposits.

### Alteration

- Some deposits have zoned alteration pipes below massive sulphides.
- Stratabound alteration zones are commonly developed in the footwall and extend along strike for 2–6 km.
- Hangingwall alteration is weakly developed or non-existent.
- Common alteration zonation toward ore is ser→ser–py→ser–chl–py→chl–py→qtz–py.

### Deposit geochemical criteria

- alteration halo is defined by Na<sub>2</sub>O depletion and MgO enrichment.
- Alteration index  $[AI = 100(MgO + K_2O)/(Na_2O + CaO + MgO + K_2O)]$  is vector toward ore (AI = 30–100).
- Ore  $100Zn/(Zn+Pb) = 60$  to 80.
- As, Sb, Hg, Tl as trace elements in Zn ores.
- Bi, Te, Mo, Co as trace elements in Cu ores.
- $\delta^{34}S$  sulphides have narrow range, commonly 0 to 15‰.
- Mean  $\delta^{34}S$  value is 17‰ less than contemporaneous SW sulphate.
- Pb isotopes form tight cluster for individual deposits that may lie on growth curve or reflect heterogeneous U/Th/Pb source area ratios.

- $\delta^{18}\text{O}$  depletion haloes approaching orebodies.
- $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  indicate majority of ore fluid is seawater.

### Surficial geochemical criteria

- Stream sediment anomalies important in discovery of Que River, Woodlawn and Wilga.
- Most deposits have strong Pb soil anomalies.
- Zn and Cu show dispersed soil anomalies.
- Gossans present at Mt Lyell, Mt Morgan, Rosebery, Woodlawn, Thalanga, Balcooma, Mt Chalmers.
- Gossan trace elements: Au, Se, Te, As, Sb, Bi, Cd, In, Tl, Hg, Sn and Ba.

### Geophysical criteria

- Regional magnetics define major volcanic units, structures and alteration.
- Regional gravity and magnetics may define position of related magnetite-series granites.
- Ores have no magnetic signature.
- EM important in discovery of Que River, Hellyer, Wilga.
- Most deposits have strong EM responses.
- Cu-rich ores have best EM response
- Zn-rich, Cu-poor ores have a very weak or non-existent EM response.
- IP defines ore zone and pyritic alteration halo.

### Fluid chemistry and source

- Ore fluids 200–350°C, 2–10 wt% NaCl
- pH 3–5.5.
- Reduced conditions in ore fluid,  $\text{H}_2\text{S} > \text{SO}_4$ .
- Metals precipitation by cooling and mixing with sea water.
- Source of fluid: seawater  $\pm$  magmatic input.
- Source of metal: Zn, Pb, Ag leached from volcanics; Cu, Au may be magmatic or leached.
- Zonation in deposits is caused by zone-refining process as hot fluids pass up through the sulphide mound.

### Comments on genesis

- Comparison with black smoker systems substantiates previous conclusions that most VHMS deposits are synvolcanic seafloor deposits.
- Seawater depths of 800–4000 m are necessary to form deposits.
- Models for seafloor sulphide formation:
  - sulphide mounds develop above hydrothermal vents and grow by upward zone refining of metals;
  - sulphide sheets or lenses may form in seafloor brine pools adjacent to vents;
  - synvolcanic sub-seafloor replacement to form stratiform massive sulphides.
- Stringer zones and sulphide pipes form by replacement and/or vein-fill processes in sub-seafloor hydrothermal vent.
- A less-favoured model invokes syntectonic replacement unrelated to volcanism.




Figure 3. Possible sources of metals for VHMS deposits.

## Introduction

Eastern Australian Palaeozoic volcanic-hosted massive sulphide (VHMS) deposits have been a major source of Cu ± Au (e.g. Mt Lyell and Mt Morgan) and a significant source of Pb–Zn–Ag (e.g. Rosebery, Hellyer, Woodlawn and Thalanga) over the last one hundred years. There is a wide range in variability or style of Australian Palaeozoic VHMS deposits, including mounds, pipes, sheets, layered deposits, stacked deposits, stockwork and disseminated deposits, and cyclic layered deposits. Although these various styles show a range of features, there is a consistent theme which strongly suggests that they all belong to the one genetic group of ore deposits.

## Distribution and age

Most VHMS deposits in Australia (24 of 30) occur within the Palaeozoic volcanic belts of eastern Australia (Table 1). The major VHMS districts are (1) the Mt Read Volcanics and correlates (Cambro-Ordovician) in Western Tasmania, host to Hellyer, Que River, Rosebery, Hercules and the Mt Lyell deposits; (2) Mt Windsor Volcanics and correlates (Cambro-Ordovician) in Queensland, host to Thalanga, Liontown, Reward, Balcooma and Dry River South, (3) the Goulburn to Cooma volcanic belt (Silurian) in southeastern New South Wales, host to the Woodlawn and Captains Flat deposits, and (4) the Enano Group volcanic sequence (Silurian) in the Benambra district of north-eastern Victoria, host to the Wilga and Currawong deposits. In addition to these four volcanic belts, individual deposits occur in the Rockhampton district of Queensland, hosted by the Devonian Capella Creek Beds (Mt Morgan deposit), the Permian Berserker Beds (Mt Chalmers deposit), and the Permian mafic volcanics (Devlin Creek deposit).

## Metal content and classification

A classification of VHMS deposits, based on the Cu ratio  $100\text{Cu}/(\text{Cu}+\text{Zn})$  and the Zn ratio  $100\text{Zn}/(\text{Zn}+\text{Pb})$ , was devised by Solomon (1976). Australian VHMS deposits fall into the following groups:

- (1) Cu deposits (Cu ratio >60, Zn ratio >60)—the Mt Lyell deposits, Mt Morgan, Mt Chalmers, Balcooma, Mt Ararat and Reward;
- (2) Zn–Cu deposits (Cu ratio <60, Zn ratio 90 to 100)—Wilga, Currawong and Devlin Creek;
- (3) Zn–Pb–Cu deposits (Cu ratio <60, Zn ratio 60 to 90)—Rosebery, Hercules, Que River, Hellyer, Thalanga, Liontown, Dry River South, Woodlawn and Captains Flat.

The Zn–Cu-type deposits are restricted to the Silurian, whereas the Cu type and Zn–Pb–Cu type occur sporadically throughout the time span from Cambrian to Permian.

## Tonnage and grade

Australian Palaeozoic deposits vary in size from 1 Mt to >90 Mt (Table 1). The most common size range is 2–5 Mt for an individual deposit, but nine deposits exceed that range: Prince Lyell (90 Mt), Mt Morgan (50 Mt), Rosebery (19.4 Mt), Woodlawn (17.7 Mt), Hellyer (17 Mt), Currawong (9.5 Mt), Thalanga (6.8 Mt), Lyell Blow (5.6 Mt) and North Lyell (5.1 Mt) (Large 1992). The polymetallic Zn–Pb–Cu deposits display the best overall grade, with an average of 11.8% Zn, 4.7% Pb, 1.0% Cu, 117 ppm Ag, 2.0 ppm Au. By comparison, the Cu-type deposits average much lower grades of 0.2% Zn, 0.0% Pb, 1.3% Cu, 8 ppm Ag, 1.6 ppm Au.

## Nature of volcanic districts

Deposits occur within the submarine portion of volcanic belts that consist of a series of complex volcanic centres with related volcanoclastic and epiclastic facies. Important aspects are:

- The volcanic package enclosing the VHMS deposits is

dominated by rhyolite (usually 60–80 per cent of pile) with lesser andesite, dacite, basalt and sediment.

- The mineralised volcanic suite consists of calc-alkaline volcanics.
- The major VHMS-bearing belts have a thick pile of rhyolitic volcanics in the base of the volcanic succession, including lavas, volcanoclastics and subvolcanic intrusives. The rhyolite sequence, which may vary in stratigraphic thickness from 1 to 3 km, either overlies a sedimentary package or its base is obscured by younger intrusive granites.
- Recent studies on physical volcanic architecture of the VHMS districts indicate that the volcanoclastic and epiclastic sequences include porous mass flows, pumiceous sandstone, siltstone, breccia and turbidites (Allen 1992, McPhie & Allen 1992). Peperitic units are also encountered at Mt Morgan and Mt Windsor (Messenger 1994, Doyle 1994).
- A mixed sequence of sediments and volcanics commonly overlies the rhyolite sequence and forms the upper part of the volcanic pile. The volcanics may be bimodal (e.g. rhyolite–andesite at Mt Chalmers), but more commonly exhibit a polymodal character (e.g. dacite, andesite, rhyolite and basalt in the Trooper Creek Formation of the Mt Windsor Volcanic pile).
- Massive sulphide deposits are generally located toward the base of the polymodal volcanic–sedimentary sequence or at the top contact of the basal rhyolite pile.
- In the Mt Read and Mt Windsor volcanic belts there are at least two important stratigraphic levels of mineralisation. In other belts only one mineralised horizon has been confidently recognised.

## Form and shape of deposits

Although a complete spectrum of shapes and forms is exhibited by the Australian deposits (Large 1992), it is possible to group them into three main types:

- **Lens and blanket deposits:** low aspect ratio with dominant zinc-rich massive sulphide lens and subordinate stringer zone. Examples are Rosebery, Thalanga, Liontown, Dry River South, Woodlawn. Generally these deposits are of the Zn–Pb–Cu and the Zn–Cu type.
- **Mound deposits:** high aspect ratio, narrow and elongate massive sulphide with a well-developed stringer zone and alteration system, directly beneath the mound. Hellyer (Zn–Pb–Cu type) and Mt Chalmers (Cu type) are good examples.
- **Pipe and stringer deposits:** cross-cutting massive sulphide pyrite–chalcopyrite pipe or stringer zones with little or no stratiform Zn-rich sulphide lenses. Examples are Prince Lyell, Lyell Blow, Highway, Mt Morgan, which all are Cu type and contain significant Au credits.

The regional distribution and the shape of the massive sulphide ores have been related to fluid focusing along syn-volcanic growth faults. In the Mt Read volcanic belt the Mt Lyell, Hercules and Rosebery deposits are located adjacent to structures representing major rift faults associated with the development of the volcanic arc (e.g. Large 1990). On a local scale, syn-volcanic growth faults have controlled fluid movement and the location and shape of deposits such as Hellyer (McArthur 1989), Mt Chalmers (Large & Both 1980), Thalanga and Reward (Gregory et al. 1990, Berry et al. 1992) and Mt Morgan (Taube 1986).

## Ore and gangue mineralogy

Considering the diverse morphology and metal ratios, Australian VHMS deposits contain a restricted range of major and minor opaque minerals. In addition to the major phases of pyrite, chalcopyrite, sphalerite and galena, ubiquitous minor phases

**Table 1. Total resource, tonnage and grades data for Australian Palaeozoic VHMS deposits over 1 Mt in size (after Large 1992 and Beams 1995).**

<i>Deposit</i>	<i>Age</i>	<i>Size</i> (Mt)	<i>Cu</i> (%)	<i>Pb</i> (%)	<i>Zn</i> (%)	<i>Au</i> (ppm)	<i>Ag</i> (ppm)	<i>Status</i>
<b>Cu type</b>								
Prince Lyell	Cambrian	90.0	0.9	nr	nr	0.3	2.0	Current mine
Mt Morgan	Devonian	50.0	0.7	nr	0.1	4.7	6.0	Past producer
Lyell Blow	Cambrian	5.6	1.3	nr	nr	2.0	61.0	Past producer
North Lyell	Cambrian	5.1	5.3	nr	nr	0.4	33.0	Past producer
Mt Chalmers	Permian	3.6	1.8	0.2	1.0	2.0	15.0	Past producer
Lyell Comstock	Cambrian	2.4	2.4	nr	nr	0.7	5.2	Past producer
Balcooma	Cambrian	2.1	3.3	nr	nr	0.5	19.0	Prospect
Crown Lyell III	Cambrian	2.1	1.4	nr	nr	0.3	4.1	Past producer
Royal Tharsis	Cambrian	1.5	1.5	nr	nr	0.5	2.8	Past producer
Cape Horn	Cambrian	1.4	1.4	nr	nr	0.4	3.3	Past producer
Highway	Cambrian	1.2	5.5	nr	nr	1.2	6.5	Current mine
Crown Lyell OC	Cambrian	1.1	1.4	nr	nr	0.5	3.9	Past producer
Mt Ararat	Cambrian	1.0	2.7	nr	nr	0.6	9.0	Prospect
<b>Zn-Cu type</b>								
Currawong	Silurian	9.5	1.7	0.9	4.3	1.3	38.0	Prospect
Wilga	Silurian	4.0	3.0	0.4	6.2	0.5	23.0	Past producer
<b>Zn-Pb-Cu type</b>								
Rosebery	Cambrian	19.4	0.7	5.0	16.2	2.9	155.0	Current mine
Woodlawn	Silurian	17.7	1.7	3.8	9.9	1.4	80.0	Past producer
Hellyer	Cambrian	17.0	0.3	7.0	13.0	2.3	160.0	Current mine
Thalanga	Cambrian	6.8	2.2	3.9	12.4	0.6	99.0	Past producer
Captains Flat	Silurian	4.2	0.7	6.0	10.0	1.7	55.0	Past producer
Dry River south	Cambrian	3.4	1.0	3.6	10.1	0.7	77.0	Prospect
Que River	Cambrian	3.1	0.6	7.5	13.5	3.4	200.0	Past producer
Hercules	Cambrian	2.6	0.4	5.2	16.7	2.7	159.0	Current mine
Liontown	Cambrian	2.0	0.5	2.3	6.6	0.9	50.0	Prospect

nr—not recorded (generally known to be less than 0.1% in the case of Pb)

are tetrahedrite, arsenopyrite, bismuth sulphosalts, cassiterite and electrum. Magnetite and pyrrhotite occur as major minerals in some Archaean Zn-Cu-type deposits (e.g. Scuddles, Gossan Hill), but are present only as minor minerals (<1 wt%) or are completely absent from the Palaeozoic VHMS deposits. Cu deposits commonly contain molybdenite, bismuthinite, various bismuth sulphosalts, bismuth tellurides and gold tellurides. Zn-Pb-Cu deposits are enriched in arsenopyrite, tetrahedrite, tennantite, boulangerite and bournonite. The dominant non-sulphide/oxide gangue minerals are sericite, chlorite, talc, barite, quartz, and various carbonates. Barite is a common constituent in the upper parts of the massive sulphide lenses of all the Zn-Pb-Cu deposits. Barite is also a common gangue mineral in the Cu-type deposits, where it may be relatively abundant in the upper parts of the Cu-rich pipe and later Zn-Pb lenses (e.g. Mt Chalmers, Large & Both 1980; Reward, Beams et al. 1989). Barite is rare in the Zn-Cu group (e.g. Wilga).

### Metal zonation

The upward-zoning sequence from Fe to Fe-Cu to Cu-Pb-Zn to Pb-Zn-Ba is now universally accepted for VHMS deposits

(e.g. Sangster 1972, Sato 1972, Large 1977, Solomon & Walshe 1979, Franklin et al. 1981, Eldridge et al. 1983, Huston & Large 1989). Mound-shaped deposits, such as Mt Chalmers and Hellyer, and the thick lenses appear to exhibit the most distinct metal zonation pattern, whereas the blanket and sheet deposits (e.g. Rosebery, Woodlawn, Wilga) show more complex zonation patterns. The Cu-rich pipes and stringer-dominated deposits (e.g. Prince Lyell, Mt Morgan) are poorly zoned.

### Alteration

Although the degree, extent, shape and zoning of hydrothermal alteration associated with Australian VHMS deposits are quite variable, there are some distinct patterns which are important for both their genetic and exploration implications. The most significant aspects are that:

- Stratabound or semi-conformable footwall alteration is more common than alteration pipes.
- Alteration pipes, especially those with significant Cu stringer mineralisation, have a central zone of intense chlorite alteration. In some cases silica alteration overprints the chlorite alteration immediately below the thickest

development of massive sulphide (e.g. Hellyer). The sericite/chlorite ratio generally increases outwards so that the chlorite pipe is surrounded by a sericite-rich envelope. Patchy carbonate alteration may extend from the outer edge of the chlorite zone through the sericite zone (e.g. Rosebery, Hellyer, Mt Chalmers).

- Where alteration pipes are not developed, chlorite may be concentrated in restricted conformable zones below Cu-rich lenses (e.g. Rosebery) or as a blanket of alteration below the full extent of the massive sulphide (e.g. Woodlawn).
- Major conformable alteration zones occur in the immediate footwall of most orebodies and are dominated by quartz-sericite  $\pm$  pyrite alteration with patchy development of chlorite and carbonate. These conformable zones extend 50–200 m into the footwall volcanic package and may extend 2–6 km along strike from the orebody.
- Hangingwall alteration is of very low intensity compared to footwall alteration, is commonly lacking in sulphides, and is best developed above the thickest portions of the massive sulphide body. Weakly developed sericite-chlorite-quartz alteration is the most common form and may be developed 10–200 m above the ore position (e.g. Mt Chalmers, Woodlawn, Rosebery). Pipes or plumes of hangingwall alteration (e.g. Hellyer) may be more common than previously recognised.

A number of studies on the chemistry of hydrothermal alteration associated with Australian VHMS deposits have been undertaken (e.g. Petersen & Lambert 1979, Large & Both 1980, Green et al. 1981, Jack 1989, Gregory et al. 1990, Gemmill & Large 1992, McGoldrick & Large 1992). The consistent geochemical patterns are:

- (1) Sodium depletion is the most common feature of footwall alteration.  $\text{Na}_2\text{O}$  content generally decreases from 2–4% in unaltered volcanics to <0.5% in the outer sericite-quartz alteration zone and commonly <0.2% in the chlorite-rich alteration.
- (2) The extensive stratabound zones of footwall sericite-quartz  $\pm$  chlorite alteration are characterised by depletion of Na, Ca, and Sr and sporadic enrichment in K, S, Ba and Sc.
- (3) Chlorite alteration as pipes or stratabound zones is also characterised by depletion of Na, Ca and Sr and by enrichment in Mg, Fe, Mn and, possibly, S, Ba, Ti and Sc.
- (4) At Hellyer, Gemmill & Large (1992) showed that the best chemical indicators of proximity to the centre of the alteration system are alteration index  $[(\text{MgO}+\text{K}_2\text{O})/(\text{Na}_2\text{O}+\text{CaO}+\text{MgO}+\text{K}_2\text{O}) \times 100]$  (Ishikawa et al. 1976), which increases from 36 to 91, S (increases from 2 to 8%), Ba (increases from 700 to 2300 ppm) and Sr (decreases from 300 to 20 ppm).
- (5) Sericite- and chlorite-bearing hangingwall alteration shows the same chemical trends as the footwall zones, but to a much lesser degree. At Mt Chalmers,  $\text{Na}_2\text{O}$  decreases, while MgO and alteration index increase as the ore horizon is approached from the hangingwall (Large 1992).
- (6) REE patterns of altered volcanics may be changed in both footwall and hangingwall alteration systems associated with VHMS deposits (Whitford et al. 1988). The most pronounced changes associated with alteration are the development of negative Eu anomalies and low Zr/Y ratios compared to the primary unaltered volcanics.

## Genesis of sulphide mound styles

The model for formation of the classic mound-shaped deposits, which can be applied to Australian examples such as Hellyer, Que River and Mt Chalmers, is based largely on the previous studies by Large (1977), Eldridge et al. (1983), Campbell et al. (1984), Lydon (1988) and Huston & Large (1989). An important

aspect of the model is the temperature evolution through time of the hydrothermal system, demonstrated at the Hellyer deposit (Khin Zaw et al. 1996) and the Kuroko deposits (Pisutha-Arnond & Ohmoto 1983), from an early intensifying stage, through a thermal maximum, finishing with a temperature decline. This thermal regime has a critical control on base and precious metal transport and deposition, owing to a strong temperature-solubility relationship for the major ore minerals.

The mound deposits develop from deposition of metal sulphides on the seafloor immediately around the hydrothermal vent. Growth of the mound occurs by upward replacement of sulphide assemblages stable at higher fluid temperatures, leading to zone refining, and deposition of the less soluble Cu minerals in the stringer zone and base of the mound, with more soluble Pb, Zn, Ag and Au minerals in the outer and upper parts of the mound (Large 1992). Immature mounds are commonly small, Pb-Zn-rich and poorly zoned (e.g. Hercules) while mature mound deposits are larger and well zoned with a complete spectrum from Cu-rich ores through to Pb-Zn and Au-Ag-rich massive sulphide (e.g. Hellyer). However, an alternative hypothesis, the brine pool model, has been postulated for the Hellyer deposit by Solomon & Khin Zaw (1997).

## Factors which control variations in VHMS deposits

The spectrum of deposit styles and the reason for their departure from the classic mound style are due to variations in a number of factors relating to the chemistry of the ore fluid, nature of the volcanic pile, and the seafloor environment. These factors include the salinity of the ore fluid, the permeability of the footwall volcanics, composition of the footwall volcanics, the temperature of the ore fluid, the  $f\text{O}_2$  and  $\text{aH}_2\text{S}$  of the fluid, and the depth of seawater.

### *Salinity variation and the brine pool model*

Sato (1972) suggested that extensive sheet-style deposits may develop from either (a) high-salinity, low-temperature fluids ( $120^\circ\text{C}$ ,  $>3\text{m NaCl}$ ), which, on venting, form a dense brine that migrates downslope to accumulate in a brine pool, or (b) high-temperature, lower salinity fluids ( $200^\circ\text{C}$ ,  $2\text{m NaCl}$ ), which, on venting, remain buoyant, resulting in dispersal of fine sulphides in the seawater column, which may then accumulate to form a thin metalliferous sediment. Solomon & Walshe (1979) and McKay & Hazeldene (1987) suggested that sheet-like deposits, such as Rosebery and Woodlawn, formed from low-salinity buoyant plumes, which spread laterally in a shallow-water environment, raining down sphalerite and galena to the seafloor. However, evidence from buoyant black-smoker plumes on the modern seafloor (e.g. Scott 1987) indicates that sheet-style massive sulphide deposits are unlikely to accumulate by this process, because of the low efficiency of precipitation and the rapid dispersal of the fine sulphide particles by seawater currents. Another possible mechanism for developing sheet-style VHMS deposits within mixed sediment-volcanic host sequences, is by stratabound replacement of particular favourable sedimentary horizons adjacent to hydrothermal fissure faults.

### *Permeability of the footwall volcanics*

The nature of hydrothermal fluid discharge up through the volcanic pile and onto the seafloor is principally controlled by the permeability of the footwall volcanic package (e.g. Green 1983). Within relatively impermeable volcanic piles, dominated by massive lava flows, significant fluid flow can only be achieved along major fault structures. Under these circumstances, well-focused fault-controlled fluids give rise to well-developed and zoned alteration pipes, with related pipe-shaped, and commonly high-grade, stringer sulphide zones below the massive sulphide mound.

In permeable volcanic sequences, such as those dominated by epiclastics or submarine mass flow breccias, hydrothermal fluid

flow would be less focused (e.g. Morton & Franklin 1987) and would permeate the sub-seafloor volcanics to produce widespread stratabound footwall alteration systems overlain by sheet-style or layered-style massive sulphide deposits (e.g. Rosebery).

### **Composition of footwall volcanics**

Volcanic piles dominated by oceanic tholeiitic basalts commonly underlie Cu–Zn-rich and Pb-poor deposits (e.g. Cyprus-type deposits, such as Devlin Creek) and modern deposits along oceanic spreading centres, whereas piles dominated by rhyolite, andesite and/or sediments commonly underlie the Pb–Zn–Cu type of deposits (e.g. Solomon 1976, Hodgson & Lydon 1977).

### **Temperature control on ore styles**

Thermodynamic studies on the solubility of ore metals (e.g. Sato 1972, Large 1977, Ohmoto et al. 1983) and fluid inclusion research on VHMS deposits (Pisutha-Arnond & Ohmoto 1983, Khin Zaw et al. 1996) have clearly demonstrated that temperature has a major control on the metal distribution and metal content of the deposits. Copper-rich deposits, such as Mt Chalmers, Mt Lyell, Mt Morgan and Balcooma are likely to have formed from hydrothermal systems which maintained output temperatures of 270–350°C over a long period of the history of ore formation, compared to Zn–Pb-rich and Cu-poor deposits (e.g. Captains Flat, Rosebery, Hellyer), where ore fluid temperatures in the range 175–235°C were probably more typical of the main phase of mineralisation.

### **Depth of seawater**

Seawater depth is the primary factor which determines whether or not the rising hydrothermal fluid boils before it reaches the seafloor (Haas 1971). No fluid inclusion evidence has been found to support the concept of ore-fluid boiling in Australian VHMS systems. However, the Cu-stockwork style of deposit at Mt Lyell may have formed due to hydrothermal fluid boiling at considerable depth below a shallow submarine or possibly sub-aerial volcanic environment. An alternative explanation for extensive sub-seafloor stockwork zones, such as Prince Lyell and the Au–Ag stockwork zone at Que River, is that they formed by interaction of hot hydrothermal fluids with particular permeable, seawater-laden, volcanic horizons (Walshe & Solomon 1981, McGoldrick & Large 1992).

### **Source of sulphur and metals**

The consistent parallel variation of sulphur isotope  $\delta^{34}\text{S}$  ratios in massive sulphide deposits and coeval marine sulphate deposits, noted worldwide by Sangster (1968), is well demonstrated in the Australian deposits and implicates seawater as a major source of sulphur for these deposits. The large amount of sulphur isotope data now available on the VHMS deposits in the Mount Read Volcanics and the Mt Windsor Volcanics shows patterns that can be interpreted to indicate a combination of reduced seawater sulphate and volcanic rock sulphur sources (Green et al. 1981, Solomon et al. 1988, Gemmell & Large 1992, 1993, McGoldrick & Large 1992).

Two alternative sources have been suggested for the metals in VHMS deposits:

- Leaching from the footwall volcanic pile and related basement rocks by heated seawater convecting above a magmatic intrusive or volcanic magma chamber (e.g. Kajiwarra 1973, Spooner & Fyfe 1973, Ohmoto & Rye 1974, Solomon 1976, Large 1977, Solomon 1981, Stolz & Large 1992).
- Direct input of a magmatic volatile phase from the magmatic system that is the source of the volcanism (Urabe & Sato 1978, Henley & Thornley 1979, Sawkins 1976, Stanton 1985, 1990).

A cartoon depicting the possible sources of sulphur and metals is shown in the exploration model. The various styles of

Australian VHMS deposits may be considered to be members of a genetic spectrum of deposits which varies from those close to the magmatic source or heat engine (e.g. Mt Morgan, Mt Lyell) and, typically, Cu–Au-rich, through the gamut of Zn–Cu and Zn–Pb–Cu styles, to those in mixed volcanic–sedimentary sequences which are well removed from the volcanic/magmatic centres (e.g. Woodlawn, Rosebery, Hercules). Based on geophysical and geochemical evidence, Large et al. (1996) proposed a model for Mt Lyell which involves mixing between deeply penetrating convective seawater, and Cu–Au–P-bearing magmatic fluid released from a shallow Cambrian granite.

### **Post-depositional metamorphic and deformational effects**

The VHMS ores along with the host volcanic piles have been variably metamorphosed from prehnite–pumpellyite facies (Hellyer) to lower greenschist (Que River), through upper greenschist facies (Rosebery), to amphibolite facies (Balcooma and Dry River south). At Rosebery, the Cambrian VHMS deposit has been affected by later metasomatic processes related to Devonian granite intrusions (Khin Zaw et al. 1997, in press). The VHMS deposits are also variably deformed from shearing, faulting and thrusting (Rosebery) to tight folding (Que River). These later effects have increased the Au tenor of ores and metallurgical viability due to remobilisation and recrystallisation of ore constituents. These later effects often generate controversies as to the timing of ore formation, and have led some workers to interpret a metamorphic origin for the Rosebery deposit (Aerden 1994) and pre-, syn-, and post-depositional origin for the Woodlawn deposit (Glen et al. 1995). At Mt Morgan, Arnold & Sillitoe (1989) considered that mineralisation formed by structurally controlled replacement associated with intrusion of the Mt Morgan tonalite. However, research by Golding et al. (1993, 1994) has showed that the Mt Morgan tonalite is younger than the age of mineralisation, and that the volcanic setting of the deposit, its shape, metal content and isotopic systematics are consistent with original volcanogenic mineralisation that has been annealed by thermal events due to intrusion of the Mt Morgan tonalite.

### **Successful exploration techniques**

Before the 1960s, all VHMS discoveries in Australia were based on outcropping gossans or mineralisation. Of the ten major discoveries since the 1960s, five relied heavily on electromagnetics (Wilga, Currawong, Que River, Hellyer, Dry River South), two on geochemistry (Woodlawn, Reward) and two on geology/gossan outcrops (Thalanga, Balcooma). Modern exploration for VHMS deposits in eastern Australia commonly involves a multidisciplinary approach, using geological mapping to define volcanic facies, and structure and alteration zones accompanied by geochemistry and geophysics to define drill targets. Litho-geochemistry as a vector to ore (e.g. Large 1995) and downhole geophysics (Bishop & Lewis 1992) have become more commonly used tools in recent years.

### **Acknowledgments**

This paper has benefited from many fruitful discussions with present and past members of the VHMS research team at the CODES Key Centre and Special Research Centre, University of Tasmania. An anonymous reviewer made many helpful comments.

### **References**

- Aerden, D.G.A.M., 1994. Microstructural timing of the Rosebery massive sulphides, Tasmania: evidence for a metamorphic origin through mobilisation of disseminated base metals. *Journal of Metamorphic Geology*, 12, 505–522.
- Allen, R.L., 1992. Reconstruction of the tectonic, volcanic and

- sedimentary setting of strongly deformed Zn–Cu massive sulphide deposits at Benambra, Victoria. *Economic Geology*, 87, 825–854.
- Allen, R.L. & Barr, D.J., 1990. Benambra copper–zinc deposits. In: Hughes, F.E. (editor), *Geology of the mineral deposits of Australian and Papua New Guinea*. Australasian Institute of Mining and Metallurgy, Monograph 14, 1539–1543.
- Arnold, G.O. & Sillitoe, R.H., 1989. Mount Morgan gold–copper deposit, Queensland, Australia: evidence for an intrusion-related replacement origin. *Economic Geology*, 84, 1805–1816.
- Beams, S.D. (editor), 1995. *Mineral deposits of North Queensland: geology and geochemistry*. James Cook University, Economic Geology Research Unit, Publication 52.
- Beams, S.D., Laing, W.P. & O'Neill, D.M., 1989. The exploration history and geology of the polymetallic Reward deposit, Mt Windsor volcanic belt, North Queensland. *Proceedings of the North Queensland Gold '89 Conference*, 95–102.
- Beams, S.D., Laurie, J.P. & O'Neill, D.M., 1990. Reward polymetallic sulphide deposit. In: Hughes, F.E. (editor), *Geology of the mineral deposits of Australian and Papua New Guinea*. Australasian Institute of Mining and Metallurgy, Monograph 14, 1539–1543.
- Berry, R.F., Huston, D.L., Stolz, A.J., Hill, A.P., Beams, S.D. & Kuronen, V., 1992. The stratigraphy, structure and volcanic-hosted mineralisation of the Mt Windsor subprovince, North Queensland, Australia. *Economic Geology*, 87, 739–763.
- Bishop, J.R. & Lewis, R.J.G., 1992. Geophysical signatures of Australian volcanic-hosted massive sulphide deposits. *Economic Geology*, 87, 913–930.
- Brathwaite, R.L., 1974. The geology and origin of the Rosebery ore deposit, Tasmania. *Economic Geology*, 69, 1086–1101.
- Campbell, I.H., McDougall, T.J. & Turner, J.S., 1984. A note on fluid dynamic processes which can influence the deposition of massive sulphides. *Economic Geology*, 79, 1905–1913.
- Cas, R., 1992. Submarine volcanism—eruption styles, products and relevance to understanding the host rock successions to VMS deposits. *Economic Geology*, 87, 511–541.
- Corbett, K.D., 1992. Stratigraphic–volcanic setting of massive sulfide deposits in the Cambrian Mount Read Volcanics, Tasmania. *Economic Geology*, 87, 564–586.
- Cox, R., Ebsworth, G.B. & Forsythe, D.L., 1990. The Wilga base–precious metal deposit, Benambra, Victoria, Australia. *Terra Nova*, 2, 262–270.
- Crawford, A.J., Corbett, K.D. & Everard, J.L., 1992. Geochemistry of the Cambrian volcanic-hosted massive sulfide-rich Mount Read Volcanics, Tasmania, and some tectonic implications. *Economic Geology*, 87, 597–619.
- Davis, L.W., 1975. Captains Flat lead–zinc orebody. In: Knight, C.L. (editor), *Economic Geology of Papua New Guinea, I. Metals*. Australasian Institute of Mining and Metallurgy, Monograph 5, 694–700.
- Doyle, M.G., 1994. Facies architecture of a submarine felsic volcanic centre: Highway–Reward, Mount Windsor Volcanics, Cambro-Ordovician, Northern Queensland. In: Henderson, R.A. & Davis, B.K. (editors), *Extended conference abstracts—New developments in the geology and metallogeny: Northern Tasmanian Orogenic zone*. James Cook University, Economic Geology Research Unit, Contribution 50, 149–150.
- Eldridge, C.S., Barton, P.B.Jr. & Ohmoto, H., 1983. Mineral textures and their bearing on formation of the Kuroko orebodies. In: Ohmoto, H. & Skinner, B.J. (editors), *Kuroko and related volcanogenic massive sulphide deposits*. *Economic Geology*, Monograph 5, 241–281.
- Franklin, J.M., Sangster, D.F. & Lydon, J.W., 1981. Volcanic-associated massive sulphide deposits. In: Skinner, B.J. (editor), *Economic Geology 75th Anniversary volume*, 485–627.
- Frets, D.C. & Balde, R., 1975. Mount Morgan copper–gold deposit. In: Knight, C.L. (editor), *Economic geology of Australia and Papua New Guinea, I. Metals*. Australasian Institute of Mining and Metallurgy, Monograph 5, 779–785.
- Gemmell, J.B., 1995. Comparison of volcanic-hosted massive sulphide mineralisation in modern and ancient back arc basins: examples from southwest Pacific and Australia. *Australasian Institute of Mining and Metallurgy, Publication 9/95, Proceedings of Pacific Rim Conference 95*, 227–232.
- Gemmell, J.B. & Large, R.R., 1992. Geological and geochemical evolution of the alteration and stringer system underlying the Hellyer volcanic-hosted massive sulphide deposit, Tasmania. *Economic Geology*, 87, 620–649.
- Gemmell, J.B. & Large, R.R., 1993. Evolution of a VHMS hydrothermal system, Hellyer deposit, Tasmania, Australia: sulphur isotope evidence. *Resource Geology*, 17, 108–119.
- Glen, R.A., Walshe, J.L., Bouffler, M., Ho, T. & Dean, J.A., 1995. Syn- and post-tectonic mineralisation in the Woodlawn deposit, New South Wales, Australia. *Economic Geology*, 90, 1857–1864.
- Golding, S., Huston, D.L., Dean, J.A., Messenger, R., Jones, I.W.O., Taube, A. & White, A.H., 1993. Mount Morgan gold–copper deposit: the 1992 perspective. In: *Australasian Institute of Mining and Metallurgy, Melbourne, Centenary Conference*, 95–111.
- Golding, S., Messenger, R., Dean, J.A., Perkins, C., Huston, D.L. & White, A.H., 1994. Mount Morgan gold–copper deposit: geochemical constraints on the source of volatiles and lead and the age of mineralisation. In: Henderson, R.A. & Davis, B.K. (editors), *Extended conference abstracts—New developments in the geology and metallogeny: northern Tasmanian orogenic zone*. James Cook University, Economic Geology Research Unit, Contribution 50, 89–95.
- Green, G.R., 1983. The geological setting and formation of the Rosebery volcanic-hosted massive sulphide ore body, Tasmania. PhD thesis, University of Tasmania (unpublished).
- Green, G.R., Solomon, M. & Walshe, J.L., 1981. The formation of the volcanic-hosted massive sulphide deposit at Rosebery, Tasmania. *Economic Geology*, 76, 304–338.
- Gregory, W., Hartley, J.S. & Wills, J.A., 1990. Thalanga zinc–lead–copper–silver deposit. In: Hughes, F.E. (editor), *Geology of the mineral deposits of Australian and Papua New Guinea*. Australasian Institute of Mining and Metallurgy, Monograph 14, 1527–1537.
- Haas, J.L.Jr., 1971. The effect of salinity on the maximum thermal gradient of a hydrothermal system at hydrostatic pressures. *Economic Geology*, 60, 940–946.
- Henley, R.W. & Thornley, P., 1979. Some geothermal aspects of polymetallic massive sulphide formation. *Economic Geology*, 74, 1600–1612.
- Hodgson, C.J. & Lydon, J.W., 1977. The geological setting of volcanogenic massive sulphide deposits and active hydrothermal systems—some implications for exploration. *Canadian Mining and Metallurgy Bulletin*, 70, 95–106.
- Huston, D.L., 1990. The stratigraphic and structural setting of the Balcooma volcanogenic massive sulphide lenses, northern Queensland. *Australian Journal of Earth Sciences*, 37, 423–440.
- Huston, D.L. & Large, R.R., 1988. The distribution, mineralogy and geochemistry of gold and silver in the north-end orebody, Rosebery Mine, Tasmania. *Economic Geology*, 83, 1181–1192.
- Huston, D.L. & Large, R.R., 1989. A chemical model for the concentration of gold in volcanogenic massive sulphide deposits. *Ore Geology Reviews*, 4(3), 171–200.
- Huston, D.L., Bottrill, R.S., Creelman, R.A., Khin Zaw, Ramsden, A.R., Rand, S., Gemmell, J.B., Jablonski, W., Sie, S.H. & Large R.R., 1992. Geological and geochemical controls on

- the mineralogy and grain size of gold-bearing phases in eastern Australian volcanic-hosted massive sulphide deposits. *Economic Geology*, 87, 542–563.
- Huston, D.L., Taylor, T., Fabray, J. & Patterson, D.J., 1992. A comparison of the geology and mineralisation of the Balcooma and Dry River South volcanic massive sulphide deposits, northern Queensland. *Economic Geology*, 87, 785–811.
- Huston, D.L., Jablonski, W. & Sie, S.H., 1996. The distribution and mineral hosts of silver in eastern Australian volcanogenic massive sulfide deposits. *Canadian Mineralogist*, 34, 529–546.
- Ishikawa, Y., Sawaguchi, T., Iwaya, S. & Horiuchi, M., 1976. Delineation of prospecting targets for Kuroko deposits based on modes of volcanism of underlying dacite and alteration halos. *Mining Geology*, 26, 105–117 (in Japanese with English abstract).
- Jack, D.J.J., 1989. Hellyer host rock alteration. MSc thesis, University of Tasmania, Hobart, 181 (unpublished).
- Kajiwara, U., 1973. A simulation of the Kuroko type mineralisation in Japan. *Geochemical Journal*, 6, 193–209.
- Khin Zaw & Large, R.R., 1992. The precious metal-rich South Hercules mineralisation, western Tasmania: a possible subsea-floor replacement volcanic-hosted massive sulfide deposit. *Economic Geology*, 87, 931–952.
- Khin Zaw, Gemmell, J.B., Large, R.R., Mernagh, T. & Ryan, C.G., 1996. Evolution and source of ore fluids in the stringer system, Hellyer VHMS deposit, Tasmania, Australia: evidence from fluid inclusion microthermometry and geochemistry. *Ore Geology Reviews*, 10, 251–278.
- Khin Zaw, Large, R.R. & Huston, D.L., 1997. Petrological and geochemical significance of a Devonian replacement zone in the Cambrian Rosebery VHMS deposit, western Tasmania. *Canadian Mineralogist*, 35, 1325–1349.
- Khin Zaw, Huston, D.L. & Large, R.R., in press. A chemical model for remobilisation of ore constituents during Devonian replacement process in the Cambrian VHMS Rosebery deposit, western Tasmania. *Economic Geology*.
- Large, R.R., 1977. Chemical evolution and zonation of massive sulphide deposits in volcanic terrains. *Economic Geology*, 72, 549–572.
- Large, R.R., 1990. The gold-rich seafloor massive sulphide deposits of Tasmania. *Geologische Rundschau*, 79(2), 265–278.
- Large, R.R., 1992. Australian volcanic-hosted massive sulfide deposits: features, styles and genetic models. *Economic Geology*, 87, 471–510.
- Large, R.R., 1995. Overview of the litho-geochemical architecture of volcanic-hosted massive sulphide deposits. 17th International Geochemical Exploration Symposium, Extended Abstracts. James Cook University, Economic Geology Research Unit, Contribution 54, 23–26.
- Large, R.R. & Both, R.A., 1980. The volcanogenic ores at Mt Chalmers, eastern Queensland. *Economic Geology*, 75, 1010–1021.
- Large, R.R. & Gemmell, J.B., 1993. Zinc-rich massive sulphide deposits in Tasmania. Australian Institute of Mining and Metallurgy, Melbourne, Publication 7/93, 153–160.
- Large, R.R., McGoldrick, J., Berry, R.F. & Young, C.H., 1988. A tightly folded, gold-rich, massive sulphide deposit: Que River mine, Tasmania. *Economic Geology*, 83, 681–693.
- Large, R.R., Huston, D., McGoldrick, P., Ruxton, P. & McArthur, G., 1990. Gold distribution and genesis in Australian volcanogenic massive sulphide deposits, and significance for gold transport models. *Economic Geology*, Monograph 5, 512–528.
- Large, R.R., Doyle, M., Raymond, O., Cooke, D., Jones, A. & Heesman, L., 1996. Evaluation of the role of Cambrian granites in the genesis of world class VHMS deposits in Tasmania. *Ore Geology Reviews*, 10, 215–230.
- Lydon, J.W., 1984. Volcanogenic massive sulphide deposits. 1: A descriptive model. *Geoscience Canada*, 11, 195–202.
- Lydon, J.W., 1988. Ore deposits models #14, volcanogenic massive sulphide deposits. 2: Genetic models. *Geoscience Canada*, 15(1), 43–65.
- Malone, E.J., Olgers, F., Cucchi, F.G., Nicholas, T. & McKay, W.J., 1975. Woodlawn copper–lead–zinc deposit. In: Knight, C.L. (editor), *Economic geology of Australia and Papua New Guinea*. 1, Metals. Australasian Institute of Mining and Metallurgy, Melbourne, Monograph 5, 701–710.
- McArthur, G.J., 1989. Hellyer. In: Burrett, C.F. & Martin, E.L. (editors), *Geology and mineral resources of Tasmania*. Geological Society of Australia, Special Publication no. 15, 144–148.
- McArthur, G.J., 1990. Metal and textural zonation in the Hellyer massive sulphide deposit. Geological Society of Australia, Special Publication no. 25, 108–109.
- McArthur, G.J. & Dronseika, E., 1990. Que River and Hellyer zinc–lead deposits. In: Hughes, F.E. (editor), *Geology of the mineral deposits of Australian and Papua New Guinea*. Australasian Institute of Mining and Metallurgy, Monograph 14, 1331–1339.
- McGoldrick, J. & Large, R.R., 1992. Geologic and geochemical controls on gold-rich stringer mineralisation in the Que River deposit, Tasmania. *Economic Geology*, 87, 667–685.
- McKay, W.J. & Hazeldene, R.K., 1987. Woodlawn Zn–Pb–Cu sulphide deposit NSW, Australia: an interpretation of ore formation from field observations and metal zoning. *Economic Geology*, 82, 141–164.
- McPhie, J. & Allen, R.L., 1992. Facies architecture of mineralised submarine volcanic sequences: Cambrian Mount Read Volcanics, western Tasmania. *Economic Geology*, 87, 587–598.
- McPhie, J. & Gemmell, J.B., 1992. Mount Read Volcanics: host sequence to Cambrian massive sulphide deposits in western Tasmania. *Bulletin of the Geological Survey, Tasmania*, 70, 161–166.
- Messenger, R., 1994. Structure and stratigraphy of the host sequence to the Mount Morgan gold–copper deposit, Queensland. 12th Australian Geological Convention, Perth, Australia. Geological Society of Australia, Abstract no. 37, 285.
- Morton, R.L. & Franklin, J.M., 1987. Two-fold classification of Archaean volcanic-associated massive sulphide deposits. *Economic Geology*, 82, 1057–1063.
- Ohmoto, H. & Rye, R.O., 1974. Hydrogen and oxygen isotopic characteristics of fluid inclusions in the Kuroko deposits, Japan. *Economic Geology*, 69, 947–953.
- Ohmoto, H., Mizukami, M., Drummond, S.E., Eldridge, C.S., Pisutha-Arnond, V. & Lenagh, T.C., 1983. Chemical processes of Kuroko formation. In: Ohmoto, H. & Skinner, B.J. (editors), *Kuroko and related volcanogenic massive sulphide deposits*. *Economic Geology*, Monograph 5, 570–604.
- Petersen, M.D. & Lambert, I.B., 1979. Mineralogical and chemical zonation around the Woodlawn Cu–Pb–Zn ore deposit, southeastern New South Wales. *Geological Society of Australia, Journal*, 26, 169–186.
- Pisutha-Arnond, V. & Ohmoto, H., 1983. Thermal history, chemical and isotopic compositions of the ore-forming fluids responsible for the Kuroko massive sulphide deposits in the Hokuroko District of Japan. In: Ohmoto, H. & Skinner, B.J. (editors), *Kuroko and related volcanogenic massive sulphide deposits*. *Economic Geology*, Monograph 5, 523–558.
- Raymond, O.L., 1996. Pyrite composition and ore genesis in the Prince Lyell copper deposit, Mt Lyell mineral field, western Tasmania, Australia. *Ore Geology Reviews*, 10, 231–250.
- Sainty, R.A., 1992. Shallow-water stratigraphy at the Mount Chalmers volcanic-hosted massive sulphide deposit, Queensland, Australia. *Economic Geology*, 87, 812–824.



- Sangster, D.F., 1968. Relative sulphur isotope abundances of ancient seas and stratabound sulphide deposits. Geological Association of Canada, Proceedings, 17, 79–91.
- Sangster, D.F., 1972. Precambrian volcanogenic massive sulphide deposits in Canada: a review. Canada Geological Survey Paper, 72-22, 44 pp.
- Sato, T., 1972. Behaviours of ore-forming solutions in seawater. Mining Geology, 22, 31–42.
- Sato, T., 1973. A chloride complex model for Kuroko mineralisation. Geochemical Journal, 7, 245–270.
- Sawkins, F.J., 1976. Massive sulphide deposits in relation to geotectonics. Geological Association of Canada, Special Paper 14, 221–240.
- Scott, S.D., 1987. Sea floor polymetallic sulphides: scientific curiosities or mines of the future. In: Teleki, P.G. et al. (editors), Proceedings NATO Advanced Research Workshop on Marine Minerals: Resource Assessment Strategies. D. Reidel Publishing Co, vol. 194: 277–300.
- Solomon, M., 1976. 'Volcanic' massive sulphide deposits and their host rocks—a review and an explanation. In: Wolf, K.H. (editor), Handbook of strata-bound and stratiform ore deposits. II, Regional studies and specific deposits. Elsevier, Amsterdam, 21–50.
- Solomon, M., 1981. An introduction to the geology and metallic mineral resources of Tasmania. Economic Geology, 76, 194–208.
- Solomon, M. & Khin Zaw, 1997. Formation on the seafloor of the Hellyer volcanogenic massive sulfide deposit. Economic Geology, 92, 686–695.
- Solomon, M. & Walshe, J.L., 1979. The formation of massive sulphide deposits on the sea floors. Economic Geology, 74, 797–813.
- Solomon, M., Eastoe, C.J., Walshe, J.L. & Green, G.R., 1988. Mineral deposits and sulphur isotope abundancies in the Mount Read Volcanics between Que River and Mount Darwin, Tasmania. Economic Geology, 83, 1307–1328.
- Spooner, E.T.C. & Fyfe, W.S., 1973. Sub-sea floor metamorphism, heat and mass transfer. Contributions to Mineralogy and Petrology, 42, 287–304.
- Stanton, R.L., 1985. Stratiform ores and geological processes. Royal Society of New South Wales, 118, 77–100.
- Stanton, R.L., 1990. Magmatic evolution and the ore type-lava type affiliations of volcanic exhalative ores. In: Hughes, F.E. (editor), Geology of the mineral deposits of Australian and Papua New Guinea. Australasian Institute of Mining and Metallurgy, Monograph 14, 101–107.
- Stolz, A.J. & Large, R.R., 1992. Evaluation of the source-rock control on precious metal grades in VMS deposits from Western Tasmania. Economic Geology, 87, 720–738.
- Taube, A., 1986. The Mount Morgan gold–copper mine and environment, Queensland: a volcanogenic massive sulphide deposit associated with penecontemporaneous faulting. Economic Geology, 81, 1332–1340.
- Urabe, T. & Sato, T., 1978. Kuroko deposits of the Kosaka mine, Northeast Honshu, Japan — products of submarine hot springs on Miocene sea floor. Economic Geology, 73, 161–179.
- Walshe, J.L. & Solomon, M., 1981. An investigation into the environment of formation of the volcanic hosted Mt Lyell copper deposits using geology, mineralogy, stable isotopes, and a six-component chlorite solid solution model. Economic Geology, 76, 246–284.
- Waters, J.C. & Wallace, D.B., 1992. Volcanology and sedimentology of the host succession to the Hellyer and Que River volcanic-hosted massive sulfide deposits, Tasmania, Australia. Economic Geology, 87, 650–666.
- Whitford, D.J., Korsch, M.J., Porritt, P.M. & Craven, S.J., 1988. Rare earth element distribution around the volcanogenic polymetallic massive sulfide deposit at Que River, Tasmania. Chemical Geology, 68, 105–119.

*Received 1 February 1996; accepted 15 October 1996*